



pADL-22c Phagemid

INSTRUCTION MANUAL

pADL™-22c Phagemid Vector for Phage Display

Catalog #: PD0110

Version: A1.11 – June 2023

Table of Contents

Description	4
Introduction	4
Content, Shipping & Storage	4
Limited Product Warranty	4
Vector Map	5
Feature Table	5
Cloning Site	6
Restriction Site Summary	6
Experimental Procedures	8
General Molecular Biology Techniques	8
Working with Filamentous Phage	8
Bacterial Strains and Helper Phage	8
Plasmid Maintenance	9
Cloning into pADL-22c	9
Sequencing of Inserts	10
Phagemid Virion Production	11
Induction Conditions & Control of Expression	11
Appendix	12
MSDS Information	12
Quality Control	12
Technical Support	12
References	12

Legal and Disclaimers

Antibody Design Labs grants to the buyer with the sale of its phage and/or phagemid vectors (the “Product”) a non-exclusive, non-transferable, royalty-free, commercial license to use Product in research conducted by the buyer (whether the buyer is an academic or a for-profit entity). The buyer is NOT granted a license to (a) use Product for human or animal therapeutic, diagnostic, or prophylactic purposes, (b) act as reseller or distributor of Product, or (c) resell, distribute, or transfer Product without modification under any name. **Antibody Design Labs** does not warrant that the use or sale of Product, the use thereof in combination with other products, or the use of Product in the operation of any process will not infringe the claims of any United States or other patent(s). If the buyer is not willing to accept the limitations of this license, without modification, buyer may refuse this license by returning Product unopened and unused. By keeping or using Product, buyer agrees to be bound by the terms of this license.

Description

Introduction

The pADL™-22c phagemid is a type 3+3 phage display vector with a cloning site for display on the N-terminal side of the full-length gene III protein. Secretion in the periplasm of the fusion protein is driven by the PelB leader peptide. A HIS tag and a HA tag followed by an amber codon are conveniently located before the gene III protein. Phage display will be done using bacterial strains that suppress the amber codon while growth on non-suppressive strains will result in the expression of free scFvs or Fab fragments in the periplasm space; a classical application of this vector is the production of free scFv and detection of its binding through the HA tag.

The pADL™-2x phagemid vector series offers optimal characteristics for phage display with optimized expression of the fusion protein for strong display, suppressible amber codon to direct the fusion either as a gene III fusion or as a secreted free entity, cloning site amenable to multiple cloning strategies and varied linker and display options. The fusion protein is under the control of the lac promoter, allowing metabolic repression by glucose and induction by IPTG. A copy of the lambda t1 terminator located downstream gene III prevents leakiness of the transcription during induction, in particular preventing excessive expression of the beta-lactamase and rapid consumption of ampicillin.

The vector contains two origins of replication, the f1 origin, which packages the single-stranded phagemid DNA into nascent virions, and the pMB1 origin of replication derived from pBR322, which results in a high-copy-number phagemid. The pMB1 sequence lacks the *rop* gene and carries a point mutation in the RNAlI transcript (G 2975 in pBR322 to T 1304 on the reverse complement strand responsible for a temperature-sensitive very high copy number phenotype (Lin-Chao 1992).

Content, Shipping & Storage

Content

VECTOR	COMPOSITION	AMOUNT
pADL™-22C	20 µl at 0.5 µg/µl of DNA vector in DNA Conservation Buffer (Tris-HCL 5 mM, EDTA 0.1 mM, pH 8.5)	10 µg

Shipping & Storage

pADL™-22c phagemid vector is shipped on wet ice. Upon receipt, store the vector at -20°C.


Limited Product Warranty

This warranty limits our liability to the replacement of this product. No other warranties of any kind express or implied, including, without limitation, implied warranties of merchantability or fitness for a particular purpose, are provided by Antibody Design Labs. Antibody Design Labs shall have no liability for any direct, indirect, consequential, or incidental damages arising out of the use, the results of use, or the inability to use this product.

For research use only; not intended for any animal or human therapeutic or diagnostic use.

Vector Map

The figure below illustrates the main features of pADL™-22c phagemid vector. The full vector sequence is available online for download in varied formats on the product web page; the total length of the vector is 3957 bp.




Feature Table

The features of pADL™-22c phagemid vector are highlighted in the following table.

FEATURE	LOCATION	DESCRIPTION
TEM1 beta-lactamase	126-986	Ampicillin resistance for selection in <i>E. coli</i> .
pMB1 origin	1141-1760	pBR322 origin for replication in <i>E. coli</i> with a high copy-number.
CAP binding site	1911-1931	Mediate the catabolite repression of the <i>lac</i> operator in the presence of glucose >1% w/v.
-35 signal	1946-1951	Lac promoter -35 signal
-10 signal	1970-1975	Lac promoter -10 signal
PelB leader sequence	2025-2087	PelB leader sequence for export in the periplasm of the host bacteria. The missing terminal methionine and alanine will have to be added during the cloning to obtain a complete leader peptide (MKYLLPTAAAGLLLLAAQPAMA) necessary for proper removal of the leader during the export process.
HIS tag	2108-2125	Peptide H H H H H H
HA tag	2132-2158	Peptide Y P Y D V P D Y A
Amber Stop codon	2162-2164	
g3p fusion coding sequence	2165-3385	Full-length gene III fusion protein coding sequence; the M13 g3p protein is fused on its N-terminal side to the HIS tag and HA tag; the exact final sequence of the fusion depends on the cloning strategy (see cloning site).
lt1	3395-3507	Lambda t1 terminator
oriF1	3628-3934	Origin of replication for phage f1

Cloning Site

Following is an illustration of pADL™-22c cloning site from the regulatory region to the g3p sequence. The PelB peptide leader sequence (translation *MKYLLPTAAAGLLLLAAQPAMA*) is interrupted by a stop codon and the two last amino acids MA will have to be re-introduced during cloning to insure proper reading frame and cleavage (on the C-terminal side of the terminal Alanine).



Restriction Site Summary

Enzyme	Site	Nb	Position	Strand	Isoschizomers
AflII	C^TTAAG	1	3381		BfrI BspTI Bst98I BstAFI MspCI Vha464I
AloI	(7/12) GAACNNNNNTCC (12/7)	1	3662		
AviII	TGC^GCA	1	688		Acc16I FspI NsbI
BamHI	G^GATCC	1	2752		
BcgI	(10/12) CGANNNNNNTGC (12/10)	1	386	-	
BfiI	ACTGGG (5/4)	1	863		BmrI BmuI
BpiI	GAAGAC (2/6)	1	3465	-	BbsI BpuAI BstV2I
BseRI	GAGGAG (10/8)	1	2540		
BsmI	GAATGC (1/-1)	1	2278		BsaMI Mva1269I PctI
BspMI	ACCTGC (4/8)	1	2790		Acc36I BfuAI BveI
BstXI	CCANNNNN^NTGG	1	2116		
BtgZI	GCGATG (10/14)	1	3706		
ClaI	AT^CGAT	1	3059		BsuTUI BspDI BseCI Bsa29I BanIII BshVI Bsu15I
DraIII	CACNNN^GTG	1	3715		AdeI
EagI	C^GGCCG	1	2047		BseX3I BstZI EclXI Eco52I
Eam1104I	CTCTTC (1/4)	1	119	-	Bst6I EarI
Eam1105I	GACNNN^NNGTC	1	908		AhdI AspEI BmeRI DriI
Eco31I	GGTCTC (1/5)	1	841		Bso31I BsaI

EcoRI	G [^] AATTC	1	2004	
FalI	(8/13)AAGNNNNNCTT (13/8)	1	3018	
GsuI	CTGGAG (16/14)	1	823	BpmI
NdeI	CA [^] TATG	1	3255	FauNDI
NheI	G [^] CTAGC	1	3386	AsuNHI BmtI BspOI
NmeAIII	GCCGAG (21/19)	1	785	-
NotI	GC [^] GGCCGC	1	2046	CciNI
PsiI	TTA [^] TAA	1	3590	AanI
PvuI	CGAT [^] CG	1	541	PleI9I BpvUI MvrI
PvuII	CAG [^] CTG	1	1856	
ScaI	AGT [^] ACT	1	430	AssI BmcAI ZrmI
SmaI	CCC [^] GGG	1	2095	Cfr9I TspMI XmaI
SpeI	A [^] CTAGT	1	2087	AhlI BcuI
AcI	AA [^] CGTT	2	310	Psp1406I
		2	683	
AlwNI	CAGNNN [^] CTG	2	1387	CaiI PstNI
		2	2719	
ApaLI	G [^] TGCAC	2	241	Alw44I VneI
		2	1487	
BaeI	(10/15)ACNNNNGTAYC (12/7)	2	2327	-
		2	2669	
BauI	CACGAG (-5/-1)	2	244	BssSI Bst2BI
		2	1628	
BciVI	GTATCC (6/5)	2	65	BfuI BsuI
		2	1592	
BglI	GCCNNNN [^] NGGC	2	2073	
		2	2094	
BseYI	CCCAGC (-5/-1)	2	1497	GsaI
		2	2074	
BspHI	T [^] CATGA	2	73	CciI PagI RcaI
		2	1081	
BsrDI	GCAATG (2/0)	2	672	Bse3DI BseMI
		2	854	
DrdI	GACNNNN [^] NNGTC	2	1693	AasI DseDI
		2	3669	
Eco57I	CTGAAG (16/14)	2	226	AcuI
		2	1274	
NaeI	GCC [^] GGC	2	2078	PdiI NgoMIV MroNI
		2	3821	
SfiI	GGCCNNNN [^] NGGCC	2	2072	
		2	2093	
TaqII	GACCGA (11/9)CACCCA (11/9)	2	195	
		2	238	
XmnI	GAANN [^] NNTTC	2	309	Asp700I MroXI PdmI
		2	3178	

Absent Sites:

AarI, AatII, AbsI, AgeI, AjuI, AlfI, ApaI, ArsI, AscI, AsuII, AvrII, BalI, BarI, BbvCI, BclI, BglII, BplI, Bpu10I, BsaBI, BsePI, BsgI, BsiWI, Bsp1407I, Bsp1720I, BssNAI, BstAPI, BstEII, Bsu36I, BtrI, CspCI, Eco47III, EcoNI, EcoRV, Esp3I, FseI, FspAI, HindIII, HpaI, KflI, KpnI, MauBI, MfeI, MluI, MreI, MroI, NarI, NcoI, NruI, OliI, PacI, PasI, PflMI, PfoI, PmaCI, PmeI, PscI, PshAI, PspXI, PsrI, PstI, RsrII, SacI, SacII, SalI, SapI, SexAI, SgfI, SgrAI, SgrDI, SnaBI, SphI, SrfI, Sse8387I, StuI, SwaI, TstI, Tth111I, XbaI, XcmI, XhoI, Zsp2I.

Experimental Procedures

General Molecular Biology Techniques

Molecular cloning and phage display should be conducted under the supervision of a qualified instructor trained to standard safety practice in a molecular biology laboratory environment. Standard molecular biology procedures can be found in a general molecular biology handbook such as Sambrook (1989).

Working with Filamentous Phage

Keep the bench clean and regularly wiped with 2% bleach to limit phage cross-contamination and only use filtered tips to prevent aerosol contaminations. Phages are known to survive standard autoclaving conditions and are not removed by 0.22 µm filtration. Phages are either killed by heat-treating dry, autoclaved materials in an oven for 4 hours at 105°C (Phage Display (2001)) or by incubation in 2% bleach for at least 1 hour. We recommend to extensively wash with hot water all glass and plastic-ware, then submerge (tubes) or incubate (flasks) with a 2% solution of bleach for at least one hour. Heat-resistant glassware can then be autoclaved in an autoclave that is never used for biological waste while sensitive plastic-ware can be used directly or at best heat-treated as described above.

Bacterial Strains and Helper Phage

Bacterial Strains

In theory, any K12 F⁺ *E. coli* strain that suppresses the amber codon is suitable for phage display using pADL-22c. Practically we recommend TG1 bacterial strain; TG1 is widely used for phage display and is well documented in the literature. TG1 suppresses amber codons and can also be made highly competent for transformation by electroporation. TG1 phenotype is highlighted below:

TG1 *supE thi-1 Δ(lac-proAB) Δ(mcrB-hsdSM)5, (r_Km_K)*
F' [traD36 proAB⁺ lacI^q lacZΔM15]

SS320 is a non-amber suppressive strain that derives from MC1061 by introduction of the F' episome (Sidhu 2000). Like most derivatives of MC1061, SS320 can be made highly competent for transformation by electroporation. SS320 can be used for example to make free scFv or free Fab secreted in the culture supernatant for performing assay in free form. SS320 phenotype is highlighted below:

SS320 *hsdR2 mcrA0 araD139 Δ(araA-leu)7697 ΔlacX74 galK16 galE15(GalS) λe14⁻ rpsL150(Str^R) spoT1 thi*
F'[proAB+lacIqlacZΔM15 Tn10 (tet^r)]

Helper Phage

We recommend CM13 helper phage. CM13 is made available by **Antibody Design Labs** under product number PH020L, which offers a highly concentrated virion preparation, eliminating the need to generate and characterize your own helper phage stocks. CM13 derived from M13KO7 by a single point mutation and produces on average twice more virions.


Plasmid Maintenance

Propagation and maintenance of pADL-22c is obtained on any *recA1*, *endA1* *E. coli* strain using LB or 2xYT medium supplemented with ampicillin 100 µg/ml as a selection marker, without glucose, and incubated at 37°C with agitation. Phagemid pADL-22c is a derivative of pBR322 with a high copy number origin of replication and usually gives high yields of plasmid DNA with most standard laboratory strains such as XL1-blue or DH5α. Some DNA stabilizing strains are known to produce smaller amounts of plasmid DNA. In case of issues, we recommend using XL10-Gold® from Agilent Technologies, Inc., on which pADL-22c plasmid DNA can be isolated in large quantities.

Cloning into pADL-22c

Primer Design and PelB Leader Sequence

A complete PelB leader sequence *MKYLPTAAAGLLLLAAQPAMA* is necessary for export in the periplasm and proper removal of the leader peptide by host proteases. In the following schema, where [NNN] represents the insert sequence and [Xxx] the translated amino acid sequence, the short hexanucleotide ATGGCN must be appended immediately to the first *BglI*/*SfiI* site to obtain a complete PelB leader encoding sequence; cleavage will occur on the C-terminal side of the terminal alanine (codon GCN).



Retention of the *SpeI* site is optional during cloning and the encoded dipeptide *ThrSer* is not known to interfere with display.

Cloning in pADL-22c Using *BglI*/*SfiI* Sites

Large libraries in the 1×10^9 range and above can easily be constructed using the double *BglI*/*SfiI* cloning site.

WORKING WITH BGL I/SFI I SITES

The *SfiI* restriction enzyme recognizes rare 8-base-long interrupted palindromes GGCCNNNN/NGGCC and leaves 3-nucleotide-long overhangs after digestion. The pADL-22c cloning site contains one *SfiI* site close to the end of the PelB leader sequence and a second *SfiI* site 8 nucleotides apart from the first site. The PelB sequence of the empty vector has an early termination by an ochre stop codon and no gene III protein is produced by the vector alone.

The *SfiI* restriction enzyme requires two copies of its recognition sequence for cleavage to occur; cleavage of the two sites happens simultaneously through interaction of two *SfiI* tetramers (Wertzell 1995). Vectors bearing two sites very close to each other are cut in trans and digestion might not complete. Therefore we strongly recommend opening pADL-22c with the alternative *BglI* restriction enzyme, which cuts the shorter 6-base-long interrupted palindromes GCCNNNN/NGGC and generates identical overhangs.

Sites open with *BglI* will re-ligate with sites open with *SfiI* as long as overhangs are complementary. Practically, the pentanucleotide NNNNN must be identical to the original vector sequence to handle both ligation of the complementary overhangs and conservation of the amino acid sequence (PelB sequence and linker to protein III). Since the overhang of the two *BglI*/*SfiI* sites are non-palindromic and different, a cut empty vector cannot ligate onto itself; it is therefore possible to follow a ligation reaction by minigel analysis since remaining unligated vector or unligated insert will migrate unchanged at their expected size.

PREPARATION OF VECTOR DNA FOR CLONING

1. On ice add successively water, *Bgl*I buffer (1x final), pADL-22c vector and *Bgl*I enzyme 5 units/μg DNA; make sure the enzyme volume does not to exceed 1/10 of the total reaction volume.
2. Incubate overnight at 37°C.
3. Inactivate for 20 min at 70°C.
4. Confirm the digestion by DNA analysis on a minigel.
5. Purify the cut vector.

For routine cloning, a standard DNA purification kit can be used directly after the digestion to remove the excess of buffer, the small DNA stuffer and leftover restriction enzyme. For library construction, phenol/chloroform extraction and/or gel purification may be required.

PREPARATION OF INSERTS

*Sfi*I digestion should be rapid and complete in 4 hours especially for fragments longer or equal to 200 bp where sites are cut in *cis*. *Bgl*I may be used when the insert sequence is known to be free of *Bgl*I site and therefore is not recommended for building antibody libraries.

Cloning using *Not*I-*Spe*I sites

The *Not*I site located in the first half of the PelB leader encoding sequence may be used in conjunction with the *Spe*I site to clone inserts. This strategy has been applied in some early phage display vectors. Consult your restriction enzyme distributor resources to identify a buffer compatible with both enzymes and follow the concentration schema given above to conduct the digestion. *Not*I and *Spe*I can be inactivated by heat before DNA purification.

Sequencing of Inserts

The following primers give both strong PCR amplification and sequencing traces. Primer locations can be found in the corresponding GenBank sequence file.

Forward or Sense Primers

phiS2 5'-ATGAAATACCTATTGCCTACGG

phiS4 5'-GCGGATAACAATTTGAATTC AAGGAGACAG

Backward, Antisense or Reverse Primers

psiR2 5'-CGTTAGTAAATGAATTTCTGTATGAGG

psiR3 5'-GCGTAACGATCTAAAGTTTGTGCG

Nested Sequencing

Often it is easier to sequence an insert by PCR on the bacterial culture supernatant or directly from a colony rather than on tediously isolated plasmids. Use the outward primers phiS2 and psiR3 together with a DNA polymerase not inhibited by bacterial cultures such as TAQ polymerase for the PCR and sequence the insert with the nested reverse primer psiR2. Use less than 1 μl of bacterial culture supernatant per 50 μl-PCR reaction or the touch of a toothpick on a colony as DNA template.

Phagemid Virion Production

A superinfection by a helper phage is necessary for phagemid pADL-22c-containing bacteria to produce virions. Please, consult the M13KO7 or CM13 helper phage manual for optimal conditions of superinfection. We recommend a rich medium such as 2xYT medium supplemented with ampicillin 100 µg/ml, kanamycin 50 µg/ml (when M13KO7 or CM13 helper phage are used), no glucose or less than 0.1% w/v, and incubation from 8 h to overnight at 30°C and 250 rpm. Supplementation with IPTG is not necessary to achieve display on the phage with pADL-22c. We recommend adding the helper phage when the bacterial culture reaches an optical density at 600 nm between 0.4 OD and 0.5 OD; large amounts of non-superinfected cells due to immunity to superinfection will decrease virion production above 0.5 OD while disparities caused by differences in phage growth rates will be amplified at a lower OD. Immunity to superinfection refers to the difficulty to transduce bacteria when protein III is expressed, as it is the case when with phagemids expressing a full-length pIII fusion protein.

Notes

- ❖ Supplementation with IPTG is not necessary during virion production but is recommended for the expression of free antibodies in the periplasm with the non-amber suppressive SS320 bacterial strain.
- ❖ Shorter incubation times 6 to 8 h long will produce less virions; we have not seen improvement of display on shorter incubation times; inversely, we have not seen sign of proteolysis of the linker after overnight incubation. Always use freshly prepared buffers from commercial concentrates during virion preparations to limit sources of proteolysis. Proteolysis usually occurs on concentrated virions; always prepare virions quickly and on ice.
- ❖ Kanamycin 50 µg/ml is enough to ensure selection with derivatives of M13KO7. Higher concentrations may be needed on very rich culture media or if your culture medium contains phosphate salts.

Induction Conditions & Control of Expression

Expression of the pIII fusion protein is under the control of the lac promoter. Because of the high copy number of the phagemid, there are not enough molecules of lacI repressor in the cell to bind to all O1/O3 operators. As a result, the lac promoter is in a induced state in TG1 and SS320. Control of the induction is done in those strains by binding of the CAP protein (catabolite activator protein) to the CAP binding site in the presence of cyclic AMP (catabolic repression). In the presence of glucose, the level of cAMP decreases, the CAP protein leaves the CAP binding site and transcription is activated. A higher level of transcription is achieved by further adding IPTG, a non-metabolizable analog of lactose.

CONDITION	STATE	NOTES
Glucose 1%	Repressed	Repression is strong but not complete. There is some leakage of the promoter. This is the recommended conditions for repression.
Glucose 2%	Repressed	A higher level of repression is achieved. Some protocols recommend this concentration (~100 mM glucose).
No Glucose	Induced	Withdrawal of the catabolic repression induces the lac promoter, resulting in expression and display.
No glucose + IPTG	Induced	A higher level of induction is obtained in the presence of IPTG (>200 µM, usually 0.5 mM). Higher levels of display may be achieved.
No Glucose + 30°C	Induced	This condition favors folding and is believed to decrease the toxicity of large proteins (e.g. Fab). This is the recommended condition for induction.

Appendix

MSDS Information

MSDSs (Material Safety Data Sheets) are available on the **Antibody Design Labs** website at the corresponding product page.

Quality Control

Specifications and quality control are detailed on the online product page. **Antibody Design Labs** certifies that the product will perform according to these specifications.

Technical Support

Visit **Antibody Design Labs'** website at www.abdesignlabs.com for technical resources, including manuals, vector maps and sequences, application notes, FAQs, etc.

FOR MORE INFORMATION OR TECHNICAL ASSISTANCE, CALL, WRITE, FAX, OR EMAIL US AT:

Antibody Design Labs

4901 Morena Blvd, Suite 203

San Diego, CA 92117

Email: support@abdesignlabs.com

Phone: 1-877-223-3104 (TOLL-FREE)

(Monday – Friday 9:00 AM – 5:00 PM PST)

References

1. LIN-CHAO S, CHEN WT, WONG TT (1992). HIGH COPY NUMBER OF THE PUC PLASMID RESULTS FROM A ROM/ROP-SUPPRESSIBLE POINT MUTATION IN RNA II. MOL MICROBIOL. 22:3385-93.
2. PHAGE DISPLAY: A LABORATORY MANUAL (2001). EDITED BY C. F. BARBAS III, D. R. BURTON, J. K. SCOTT, AND G. J. SILVERMAN. COLD SPRING HARBOR, LABORATORY PRESS, COLD SPRING HARBOR, NY.
3. SAMBROOK, J., FRITSCH, E.F., AND MANIATIS, T. (1989). IN MOLECULAR CLONING: A LABORATORY MANUAL. COLD SPRING HARBOR LABORATORY PRESS, NY, VOL. 1, 2, 3.
4. SCOTT JK & BARBAS CF (2001). PHAGE-DISPLAY VECTORS (2.1-2.19) IN PHAGE DISPLAY: A LABORATORY MANUAL. EDITED BY C. F. BARBAS III, D. R. BURTON, J. K. SCOTT, AND G. J. SILVERMAN. COLD SPRING HARBOR, LABORATORY PRESS, COLD SPRING HARBOR, NY.
5. SIDHU SS, LOWMAN HB, CUNNINGHAM BC, WELLS JA (2000). PHAGE DISPLAY FOR SELECTION OF NOVEL BINDING PEPTIDES. METHODS ENZYMOL.328:333-63.
6. WERTZELL L.M. ET AL., (1995). THE SFII RESTRICTION ENDONUCLEASE MAKES A FOUR-STRAND DNA BREAK AT TWO COPIES OF ITS RECOGNITION SEQUENCE. J. MOL. BIOL. 248:581-595

This product is subject to Antibody Design Labs Terms & Conditions of Sales available online at <http://www.abdesignlabs.com/terms/>.

© 2023 Antibody Design Labs. All rights reserved.